



Little Treasure™ White

Arabis Little Treasure™

syngenta *flowers*

Culture Guide

Botanical name: *Arabis caucasica*

Product form: Seed

Containers: Quarts, Gallons

Habit: Mounding

Vernalization: Required

Garden Specifications

Garden Height: 4–8" (10–20 cm) tall

Garden Width: 4–8" (10–20 cm) wide

Exposure: Part sun–Full sun

USDA zone: 3–7

AHS zone: 7–3

Product use: Containers

Germination

Stages 1 & 2

Germination time: 7–10 days

Media temp: 68–70 °F (19–21 °C)

Chamber: Optional

Light: Required for germination

Seed cover: No

Moisture level: 4 - WET: Soil is dark brown but not shiny, no free water is seen at the surface of the soil, when pressed or squeezed water drips easily, and trays are heavy with a visible bend in the middle.

Rec. tray size: 288-cell tray

Seeds per cell: 3–4

Young Plant Production

Stages 3 & 4

Temperature

Day: 64–66 °F (18–19 °C)

Night: 60–62 °F (16–17 °C)

Average daily temperature: 62 °F (17 °C)

Lighting

Day extension lighting: Not necessary

Light intensity: 200–250 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for the first two weeks after sowing or until root development occurs. Light levels can be increased up to 600 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ as rooting increases and the seedling matures.

Day length response: Day neutral

Daily light integral: 10–12 $\text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$

Media pH: 5.5–5.9

Media EC: SME 0.25 to 0.5 mS/cm



Little Treasure™ Deep Rose

Fertilizer: 50–75 ppm N

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 2 and 3.

2 - MEDIUM: Soil is light brown in color, no water can be extracted from soil, and soil will crumble apart.

3 - MOIST: Soil is brown in color, strongly squeezing the soil will extract a few drops of water, and trays are light with no visible bend.

Pinching: No

Plant growth regulators (PGRs): Not necessary

Plug grow time: 6–7 weeks in a 288-cell tray

Comments: Do not keep plugs overly wet and avoid high ammonium feeds to avoid lush growth.

Bulking and Vernalization

Vernalization: Required

Bulking

Bulking Time: See Scheduling Section

Temperature: 66–68 °F (19–20 °C)

Pinch: No

PGR: Not Necessary

Vernalization

Vernalization Time: 5–8 weeks

Temperature: 40–50 °F (4–10 °C)

Tech tip: When bulking, allow only enough time to grow the plant within 1" (2.5 cm) of the edge of the finish container before the start of vernalization. This will decrease the chance of disease and make growth more manageable in the spring.

Finishing

Temperature

Day: 68–70 °F (20–21 °C)

Night: 60–62 °F (16–17 °C)

Average daily temperature: 64 °F (18 °C)

Lighting

Day extension lighting: Not necessary

Light intensity: 1,200–1,600 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$

Day length response: Day neutral

Daily light integral: 14–16 $\text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$

Transplanting: Transplant directly into the finished container. Place the rooting media slightly above the level of media in the container. Make sure the root ball is covered and that the plug is situated in the center of the pot.

Media pH: 5.8–6.2

Media EC: SME EC: 0.9–1.3 mS/cm, PourThru EC: 1.4–2.0 mS/cm

Fertilizer: 75–125 ppm N

Pinching: No. Pinching is not recommended or needed.

Plant growth regulators (PGRs): If grown with low ammonium feed and cool temperatures, PGRs should not be needed. But, if needed a B-Nine® WSG spray at 2,500 ppm can be used.

Tech tip: Photoperiod has no effect on the time to flower or the number of blooms produced on Little Treasure.



Try Chrysal Alesco®, a postharvest foliar spray, to protect ethylene sensitive crops during shipping and retail.

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 2 and 3.

2 - MEDIUM: Soil is light brown in color, no water can be extracted from soil, and soil will crumble apart.

3 - MOIST: Soil is brown in color, strongly squeezing the soil will extract a few drops of water, and trays are light with no visible bend.

Common pests: Aphids

Common diseases: Botrytis, Pythium, Rhizoctonia

Scheduling

Size	Crop Time			Plants Per Pot
	Bulk	Vern.	Finish	
1.0 quart (4.5 to 5 inch)	5–6 weeks	5–8 weeks	6–7 weeks	1–2 ppp
1.25 to 2.5 quart (5.5 to 6.5 inch, trade gallon)	6–7 weeks	5–8 weeks	6–7 weeks	2–3 ppp

Estimated finish crop time is from transplant of a 288-cell tray and finished at an average daily temperature (ADT) of 64 °F (18 °C).

Example crop schedule for a 1.0 quart

Weeks From Transplant	Description
1 week	Transplant one plug in the center of a quart container
6 weeks	Finish the bulking period once the plant is 1/3 to 1/2 the size of the final container. Then, begin the vernalization period
13 weeks	Finish the vernalization period and start to increase the temperatures to force flowering
16 weeks	Finish



Little Treasure™ White