



Aubrieta Audrey™

Culture Guide

syngenta® flowers

Botanical name: *Aubrieta hybrida*

Product form: Seed

Containers: Quarts, Gallons

Habit: Mounding

Vernalization: Required

Garden Specifications

Garden Height: 6–8" (15–20 cm) tall

Garden Width: 6–8" (15–20 cm) wide

Exposure: Full sun

USDA zone: 5–9

AHS zone: 9–6

Product use: Containers, Landscapes, Combos

Germination

Stages 1 & 2

Germination time: 8–14 days

Media temp: 64–68 °F (18–20 °C)

Chamber: Optional

Light: Not required for germination

Seed cover: Yes

Moisture level: 4 - WET: Soil is dark brown but not shiny, no free water is seen at the surface of the soil, when pressed or squeezed water drips easily, and trays are heavy with a visible bend in the middle.

Rec. tray size: 288–cell tray

Seeds per cell: 2–3

Young Plant Production

Stage 3 & 4

Temperature

Day: 62–64 °F (17–18 °C)

Night: 58–60 °F (14–16 °C)

Average daily temperature: 60 °F (16 °C)

Lighting

Day extension lighting: Not necessary

Light intensity: 200–250 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for the first two weeks after sowing or until true leaf development occurs. Light levels can be increased up to 600 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ as rooting increases and the seedling matures.

Day length response: Day neutral

Daily light integral: 8–10 $\text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$

Media pH: 5.8–6.2

Media EC: SME 0.5 to 0.75 mS/cm



Audrey™ Purple Shades

Fertilizer: 50–75 ppm N

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 2 and 3.

2 - MEDIUM: Soil is light brown in color, no water can be extracted from soil, and soil will crumble apart.

3 - MOIST: Soil is brown in color, strongly squeezing the soil will extract a few drops of water, and trays are light with no visible bend.

Pinching: No

Plant growth regulators (PGRs): B-Nine® WSG spray at 1,500–2,500 ppm, or Altercel™ spray at 500–750 ppm can be used if necessary to tone plugs.

Plug grow time: 5–6 weeks in a 288–cell tray

Comments: Keep evenly moist and don't allow to dry out completely. Although germination is about 65–70%, it is best seeded with 2 seeds/plug to maintain even developing plants. Too many seeds per plug results in uneven finished pots.

Bulking and Vernalization

Vernalization: Required

Bulking

Bulking Time: See Scheduling Section

Temperature: 64–66 °F (18–19 °C)

Pinch: No

PGR: Not Necessary

Vernalization

Vernalization Time: 6–10 weeks

Temperature: 40–50 °F (5e–10 °C)

Tech tip: Before vernalization, plants should be well rooted and the foliage should fill ½ to ⅓ of the overall pot diameter. Keep fertility on the low side going into vernalization to avoid excess soft growth. This decreases the chance of disease and allows room for growth during finishing.

Finishing

Temperature

Day: 66–68 °F (19–20 °C)

Night: 58–60 °F (14–16 °C)

Average daily temperature: 64 °F (18 °C)

Lighting

Day extension lighting: Not necessary

Light intensity: 1,200–1,600 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$

Day length response: Day neutral

Daily light integral: 14–16 $\text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$

Transplanting: Transplant directly into the finished container. Place the rooting media slightly above the level of media in the container. Make sure the root ball is covered and that the plug is situated in the center of the pot.

Media pH: 5.5–5.9

Media EC: SME EC: 0.9–1.3 mS/cm, PourThru EC: 1.4–2.0 mS/cm

Fertilizer: 75–125 ppm N

Pinching: No. Pinching is not recommended or needed.

Plant growth regulators (PGRs): If required, apply B-Nine® WSG spray at 2,500–3,750 ppm, or Altercel™ spray at 500–1,000 ppm.

Tech tip: When applicable, grow cool to keep the plants compact and uniform.



Try Chrysal Alesco®, a postharvest foliar spray, to protect ethylene sensitive crops during shipping and retail.

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 2 and 3.

2 - MEDIUM: Soil is light brown in color, no water can be extracted from soil, and soil will crumble apart.

3 - MOIST: Soil is brown in color, strongly squeezing the soil will extract a few drops of water, and trays are light with no visible bend.

Common pests: Aphids

Common diseases: Botrytis, Pythium, Rhizoctonia

Scheduling

Size	Crop Time			Plants Per Pot
	Bulk	Vern.	Finish	
1.0 quart (4.5 to 5 inch)	5–6 weeks	6–10 weeks	14–18 weeks	1–2 ppp
1.25 to 2.5 quart (5.5 to 6.5 inch, trade gallon)	6–7 weeks	6–10 weeks	14–18 weeks	2–3 ppp

Estimated finish crop time is from transplant of a 288-cell tray and finished at an average daily temperature (ADT) of 64 °F (18 °C).

Example crop schedule for a 2.5 quart

Weeks From Transplant	Description
1 week	Transplant two plugs into the center of the pot
6 weeks	With bulking complete, start to lower temperatures and begin the vernalization process
12 weeks	With vernalization complete, start to raise the temperatures to 64 °F (18 °C) ADT to start to forcing flowers
22 weeks	If needed, apply PGRs to help control size
26 weeks	Finish

Audrey™ Sky Blue

